

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/256715057>

# First report of *Agrocybe retigera* (Speg.) Singer (Bolbitiaceae, Agaricales) from Brazil

Article · January 2005

CITATION

1

READS

57

## 2 authors:



Vagner G. Cortez

Universidade Federal do Paraná

89 PUBLICATIONS 443 CITATIONS

[SEE PROFILE](#)



Rosa Mara B. da Silveira

Universidade Federal do Rio Grande do Sul

79 PUBLICATIONS 503 CITATIONS

[SEE PROFILE](#)

## Some of the authors of this publication are also working on these related projects:



Monitoring of macro and microfungi diversity (systematics and ecology) in the Neotropics [View project](#)



Taxonomy and phylogeny of Fomitiporia (Hymenochaetaeaceae, Basidiomycota) in Brazil [View project](#)

## FIRST REPORT OF *Agrocybe retigera* (SPEG.) SINGER (BOLBITIACEAE, AGARICALES) FROM BRAZIL

In 1922, the Italian mycologist and botanist Carlos Spegazzini described the fungus *Naucoria retigera* Speg., which was collected by himself in Paraguay (SPEGAZZINI, 1922). Years later, Dr. Rolf Singer, during his studies on type materials described by Spegazzini (SINGER, 1950) studied this species and transferred the mushroom to the genus *Agrocybe* Fayod (family Bolbitiaceae Singer). In the same work, Singer established that *Naucoria semiorbicularis* var. *lacunosa* Murrill from Florida (USA) is a taxonomic synonym of *A. retigera*. Since then, a few records of this species are known: it was reported from Argentina (SINGER; DIGILIO, 1951), Venezuela (DELGADO; URDANETA, 2002), Hawaiian Islands (DESJARDIN et al., 2005), and Mexico (GUZMÁN, 2003).

The genus *Agrocybe* has received little attention in Brazil. The contribution of Watling (1992) from the state of Paraná is undoubtedly the most important account of the Brazilian Bolbitaceae, 11 species of *Agrocybe* are reported and/or described therein. In the revision of the Agaricales *s.l.* from the state of São Paulo, 6 *Agrocybe* species are keyed by Pegler (1997). In the above-cited works *A. retigera* is not reported, thus being unknown in Brazil.

Recently the authors collected and studied specimens of *Agrocybe*, which agree in all aspects with the description of *A. retigera* as presented by Singer (1950) and Singer and Digilio (1951). These materials are described and illustrated as follows:

*Agrocybe retigera* (Speg.) Singer, *Lilloa* 23: p. 213, 1950

Figs. 1-6

≡ *Naucoria retigera* Speg., An. Mus. Nac. Buenos Aires 31: p. 363, 1922.

= *Naucoria semiorbicularis* var. *lacunosa* Murrill, *Mycologia* 35: p. 431, 1943.

**Pileus** 17-41 mm in diameter, conic-convex, hemispheric or plano-convex, umbonate, yellow-cream colored to pale brownish in age, surface

sub-viscid to moist, rugose to sulcate (escrobiculate) in younger basidiomata, but disappearing the older ones, margin somewhat striate. **Gills** adnexed to sinuate, firstly cream-colored to gray, finally tobacco-brown when mature, membranous. **Stipe** 51-77 × 3-6 mm, central, cylindrical with a slightly swollen base, cream-colored, surface covered with thin fibrils, hollow. **Veil** absent. **Context** fleshy and soft, white. **Spore print** tobacco-brown.

**Basidiospores** 12-17 × 6.5-9 µm, ellipsoid both in side and frontal view, brownish under KOH, smooth and thick-walled, with a conspicuous and wide germ pore, apiculus present. **Basidia** 26.5-33.5 × 8-15 µm, clavate-ventricose, mainly tetrasporic, but some bear two sterigmata. **Pleurocystidia** 35-70 × 15-25 µm, mainly lageniform and ventricose, with a rounded or subcapitate apex, thin-walled, hyaline. **Cheilocystidia** 29.5-61 × 10.5-21.5 µm, utriform to ventricose, thin-walled, hyaline. **Pileipellis** cellular, composed by sphaeropedunculate, thin-walled, 30-54 × 16-30 µm, hyphae. **Gill trama** regular. **Clamp connections** present.

**Habitat:** growing in numerous groups on soil of gardens, between grasses.

**Material examined: BRAZIL. Rio Grande do Sul: Porto Alegre**, Av. Nilo Peçanha, 18.III.2005, *L.F.P. Lima* 129 (ICN); **Santa Maria**, Universidade Federal de Santa Maria, Camobi, 25.I.2002, *Cortez 003/02* (SMDB 9547); **Santa Cruz do Sul**, Centro de Radiologia, 16.II.1990, *A.B. Pereira* (HCB 15673 – as *Stropharia* sp.).

The characters used by Singer and Digilio (1951) for the delimitation of this taxon were the rugose pileus surface, absence of an annular veil, presence of large pleurocystidia and cheilocystidia, and the larger basidiospores, with a conspicuous germ-pore. All that elements are present in the collected specimens. Singer (1950) reported shorter pleurocystidia (20-45 µm) based on Murrill's materials from Florida, but did not discuss about these structures in South American collections. The synonymy is according to Singer (1950) and Watling and Gregory (1981). *Agrocybe semiorbicularis* (Bull.: St. Amans) Fayod, a European species, seems to come very close to *A. retigera*, but

the former does not possess pleurocystidia, as indicated by Singer (1950).

*Agrocybe retigera* is a ruderal and probably very common species. It has been seen growing in gardens, parks and meadows between the months of January and March. It is surprising that this species was overlooked for so many times. *Agrocybe retigera* is here reported for the first time in Brazil, and represents an extension of its distribution. As previously discussed, the species is known from Florida to Argentina, but unnoticed in several Central and South American countries where it probably occurs.

#### ACKNOWLEDGEMENTS

The authors thank to Dr. Roy Watling for kindly reading the manuscript and made invaluable suggestions. Luis F. Lima is acknowledged for collecting specimens.

#### REFERENCES

DELGADO, A. E.; URDANETA, L. M. Hongos Basidiomycota, orden Agaricales, en cinco municipios del estado Zulia, Venezuela. *Revista de la Facultad de Agronomía (LUZ)*, Caracas, v. 19, p. 56-70, 2002.

DESJARDIN, D. E.; HEMMES, D.; WONG, G. J. **Agaricales of the Hawaiian Islands**. Available at: <<http://www.mycena.sfsu.edu/pages/hawaiian/Agaricales.html>>. Access in 18 Mar. 2005.

GUZMÁN, G. **Los hongos de el Éden, Quintana Roo**: introducción a la micobiota tropical de México. Xalapa: Instituto de Ecología, A.C./CONABIO, 2003. 316 p.

PEGLER, D.N. **The agarics of São Paulo, Brazil**. London: HMSO/Royal Botanic Gardens, Kew, 1997. 68 p.

SINGER, R. Type studies on Basidiomycetes IV. *Lilloa*, Tucumán, v. 23, p. 5-461, 1950.

SINGER, R.; DIGILIO, A. P. L. Pródromo de la Flora Agaricina Argentina. *Lilloa*, Tucumán, v. 25, p. 5-461, 1951.

SPEGAZZINI, C. Fungi Paraguayensis. *Anales del Museo Nacional de Buenos Aires*, Buenos Aires, v. 31, p. 355-450, 1922.

WATLING, R. Observations on the Bolbitiaceae – 30. Some Brazilian taxa. *Boletín de la Sociedad Argentina de Botánica*, Buenos Aires, v. 28, p. 77-103, 1992.


WATLING, R.; GREGORY, N. M. **Census catalogue of world members of the Bolbitiaceae**. Vaduz: J. Cramer, 1981. 224 p. (Bibliotheca Mycologica, 82).

**Vagner Gularte Cortez<sup>1</sup>**  
**Rosa Mara Borges da Silveira<sup>2</sup>**

Recebido em: 31.03.05; aceito em: 28.09.05.

<sup>1</sup> Bolsista do CNPq. Mestrando do Programa de Pós-graduação em Botânica, Universidade Federal do Rio Grande do Sul. Av. Bento Gonçalves, 9500, CEP 91501-970, Porto Alegre, RS, Brasil – E-mail: [cortezvg@yahoo.com.br](mailto:cortezvg@yahoo.com.br)

<sup>2</sup> Departamento de Botânica, Universidade Federal do Rio Grande do Sul.



**Figs. 1-6.** *Agrocybe retigera* (Speg.) Singer: **1.** Basidiomata and pileus section. **2.** Basidia. **3.** Basidiospores. **4.** Cheilocystidia. **5.** Pleurocystidia. **6.** Pileipellis hyphae.